

(b) Find $\Delta^{n+1} x^n = ?$

(c) Write down Newton's forward interpolation formula.

(d) The Newton-Raphson method is also called as

- (i) tangent method
- (ii) secant method
- (iii) chord method
- (iv) diameter method

(Choose the correct option)

(e) In the general Quadrature formula Simpson's one third rule is obtained by putting

- (i) $n = 1$
- (ii) $n = 2$
- (iii) $n = 3$
- (iv) $n = 4$

(Choose the correct option)

(f) The value of $\int_0^{\pi/4} \frac{dx}{1+x^2}$ is

- (i) 0
- (ii) 1
- (iii) 2
- (iv) None of the above

(Choose the correct option)

(g) Where is Euler's method used ?

2. Answer the following questions : $2 \times 4 = 8$

(a) Define rate of convergence and order of convergence of a sequence.

(b) Evaluate : $\frac{\Delta^2}{E} x^3$

(c) Construct a divided difference table from the following data :

x	-1	1	2	3
y	-21	15	12	3

(d) Why is Lagrange's formula considered to be of more general nature than Newton's formula?

3. Answer **any three** questions : $5 \times 3 = 15$

(a) What do you mean by algorithm? Use the statistics algorithm to compute the mean and standard deviation of the following data : $1 + 4 = 5$

1, 3, 5, 7, 9

(b) Find a root of the equation

$$x^3 - 4x - 9 = 0$$

using the bisection method correct up to 3 decimal places.

(c) Show that

(i) $\delta \equiv \nabla(I - \nabla)^{-\frac{1}{2}}$

(ii) $E\Delta \equiv \Delta E$ $3 + 2 = 5$

(d) Find the rate of convergence of Newton-Raphson method.

(e) Using Lagrange's interpolation formula for unequal interval, find the values of $f(2)$ and $f(15)$ from the following data :

x	4	5	7	10	11	13
$f(x)$	48	100	294	900	1210	2028

4. Answer the following questions : $10 \times 3 = 30$

(a) Determine the root of

$xe^x - 2 = 0$ by the method of false position. Perform *five* iterations.

OR

Form an LU decomposition of the following matrix :

$$A = \begin{pmatrix} 1 & 4 & 3 \\ 2 & 7 & 9 \\ 5 & 8 & -2 \end{pmatrix}$$

(b) Let x_0, x_1, \dots, x_n be $(n+1)$ distinct points on $[a, b]$. If f is continuous on $[a, b]$ and has n continuous derivatives on (a, b) , then prove that there exist some $\xi \in (a, b)$ such that

$$f[x_0, x_1, \dots, x_n] = \frac{f^n(\xi)}{\xi!}$$

where $f^n(x) = \frac{d^n f(x)}{dx^n}$.

Find the interpolating polynomial from the data given below using divided differences:

$$\begin{array}{l} x : -2 \quad 0 \quad 2 \\ f(x) : 4 \quad 2 \quad 8 \end{array} \quad \begin{array}{l} 5+5=10 \end{array}$$

OR

Derive the formula for finding first and second order derivatives using Newton's forward difference formula.

Given that

X:	1.0	1.1	1.2	1.3	1.4	1.5	1.6
Y:	7.989	8.403	8.781	9.129	9.451	9.750	10.031

Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$ at $x = 1.1$ 5+5=10

(c) Define numerical integration.

Obtain a general quadrature formula for

$$\int_a^b f(x) dx.$$

Hence deduce Simpson's $\frac{1}{3}$ rd rule.

$$1+5+4=10$$

OR

Write a short note on Euler's method. Give the geometric interpretation of Euler's method.

Give an algebraic interpretation of Euler's method.

Solve by using Euler's method:

$$y' = x + y; \quad y(0) = 2 \quad \text{for } 0 \leq x \leq 1$$

$$h = 0.5$$

$$2+2+2+4=10$$

Total number of printed pages-7

3 (Sem -4/CBCS) MAT HC 2

2023

MATHEMATICS

(Honours Core)

Paper : MAT-HC-4026

(**Numerical Methods**)

Full Marks : 60

Time : Three hours

The figures in the margin indicate full marks for the questions.

1. Answer the following questions as directed :
 $1 \times 7 = 7$

(a) What is the order of convergence of Regula-Falsi method ?

(i) 2.312

(ii) 2.231

(iii) 1.618

(iv) 1.321

(Choose the correct option)

Contd.